



# 18-0-3 25% XCU® with Dimension® 0.17% Herbicide

Professional turf fertilizer with Dimension® herbicide  
PremiumTurf Grade (SGN 200)

## Product Features

- Fertilizer with Dimension® 0.17% Herbicide provides season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes, and in established landscape ornamentals.
- This product also provides postemergent crabgrass control up to the 4-leaf stage.
- 25% of the nitrogen is derived from XCU® polymer-coated, sulphur-coated urea. XCU® slow-release nitrogen provides extended feeding.

## Turf Benefits

- XCU® slow-release fertilizer uses breakthrough technology to provide a unique coating that allows gradual and consistent nutrient release delivering up to 8 weeks of continuous feeding.

## Product Characteristics

<b>Product Code:</b>	P000798
<b>Product Type:</b>	Professional Turf Fertilizer with Dimension® Herbicide
<b>Particle Size:</b>	Premium SGN 200
<b>Net Weight:</b>	50 lb
<b>Bulk Density:</b>	70 lb/ft³
<b>Coverage:</b>	14,710 ft² @ 3.4 lb/M

## Guaranteed Analysis

	%
Total Nitrogen (N)	18.0
18.0% Urea Nitrogen*	
Soluble Potash (K₂O)	3.0
Chlorine not more than (Cl)	2.2
Derived from: Urea, Polymer-coated, sulfur coated urea and Muriate of potash.	

\* 4.2% Slow release nitrogen from polymer-coated, sulfur-coated urea.

**Active Ingredient:** 0.17% Dithiopyr



XCU® is registered trademark of Agrium.

Dimension® is a Trademark of the Dow Chemical Company ('Dow') or an affiliated company of Dow.

**For more information, contact Nutrite at:**

**1-800-265-8865**

**[www.nutrite.com](http://www.nutrite.com)**



## Nutrite Professional Turf Fertilizer 18-0-3 with Dimension® 0.17% Herbicide



Further Treatment Advice.

For further information, contact 1-800-322-8644 or visit [www.nutrite.com](http://www.nutrite.com).

### ENVIRONMENTAL INFORMATION

## Nutrite Professional Turf Fertilizer 18-0-3 with Dimension® 0.17% Herbicide

### For season-long control of crabgrass and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

**NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL PURPOSES, OR FOR RESEARCH PURPOSES. USE FOR PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATION AND BEING GROWN IN ORNAMENTAL GARDENS OR PARKS, OR ON GOLF COURSES OR LAWNS.**

In the state of New York, this product may be applied only to commercial applicators at no more than 284 lb. (0.5 lb active ingredient) per acre (6.8 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

This product contains 0.085 pound of the active ingredient dimorphoate per 50 pound bag.

Dimension is a trademark of the Dow Chemical Company (Dow) or an affiliated company of Dow.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

EPA REG. NO. 62771-9-485-87245

EPA EST. NO. 87245-LAN-001

**Product Code: P000798**

**Net Wt. 50 lb (22.68 kg)**

**MANUFACTURED, QUARANTEED AND DISTRIBUTED BY:**

Ferti Technologies  
560 Rhéaume P.O. Box 129  
Saint-Michel, QC  
Canada, J0L 2J0

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

##### Causes Eye Irritation • Harmful If Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after use. If eye irritation continues, seek medical advice or treatment.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### Application Directions for Turf

For improved weed control, this product is limited to crabgrass. Postemergence application is recommended for crabgrass. For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Improved Weed Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Applications

When applied to preemergence control of crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Applications

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Spring Pre-emergence Control

When applied to crabgrass in the spring, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by rainfall or irrigation.

If eye irritation continues, seek medical advice or treatment.

Use of split (half-rate) applications delayed 5-10 weeks, again may provide improved weed control.

Do not use clippings from treated turf for mulching around vegetables or fruit trees.

Do not graze livestock or pets enter treated areas if settled.

##### For Fall Pre-emergence Control

When applied to crabgrass in the fall, this product is limited to crabgrass.

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

The herbicide and fertilizer/ammonium in this product are not effective until activated by